

Newborn Screening ACT Sheet

[Increased Phenylalanine]

Phenylketonuria (PKU)

Differential Diagnosis: Phenylketonuria (Classical PKU); non-PKU mild hyperphenylalaninemia; pterin defects; transient hyperphenylalaninemia.

Condition Description: In PKU the phenylalanine from ingested protein cannot be metabolized to tyrosine because of deficient liver phenylalanine hydroxylase (PAH). This causes elevated phenylalanine. Pterin defects result from deficiency of tetrahydrobiopterin (BH4), the cofactor for PAH and other hydroxylases. This produces not only increased phenylalanine but also neurotransmitter deficiencies.

YOU SHOULD TAKE THE FOLLOWING ACTIONS IMMEDIATELY:

- Contact family immediately to inform them of the newborn screening result.
 - Consult with pediatric metabolic specialist.
 - Evaluate the newborn and refer as appropriate.
 - Initiate confirmatory/diagnostic tests in consultation with metabolic specialist.
 - Provide the family with basic information about PKU and dietary management.
 - Report findings to newborn screening program.
-

Diagnostic Evaluation: Plasma amino acid analysis which shows increased phenylalanine without increased tyrosine (increased phenylalanine:tyrosine ratio). Urine pterin analysis and red blood cell DHPR assay will identify pterin defects. Consider PAH mutation testing.

Clinical Considerations: Asymptomatic in the neonate. If untreated PKU will cause irreversible mental retardation, hyperactivity, autistic-like features, and seizures. Treatment will usually prevent these symptoms. Pterin defects cause early severe neurologic disease (developmental delay/seizures) and require specific therapy.

Additional Information:

[Gene Reviews](#)

[Genetics Home Reference](#)

[PKU](#)

[Tetrahydrobiopterin Deficiency](#)

Referral (local, state, regional and national):

[Testing](#)

[Clinical Services](#)

[Find Genetic Services](#)

Disclaimer: This guideline is designed primarily as an educational resource for clinicians to help them provide quality medical care. It should not be considered inclusive of all proper procedures and tests or exclusive of other procedures and tests that are reasonably directed to obtaining the same results. Adherence to this guideline does not necessarily ensure a successful medical outcome. In determining the propriety of any specific procedure or test, the clinician should apply his or her own professional judgment to the specific clinical circumstances presented by the individual patient or specimen. Clinicians are encouraged to document the reasons for the use of a particular procedure or test, whether or not it is in conformance with this guideline. Clinicians also are advised to take notice of the date this guideline was adopted, and to consider other medical and scientific information that become available after that date.

LOCAL RESOURCES: Insert State newborn screening program web site links

State Resource site (insert state newborn screening program website information)

Name	
URL	
Comments	

Local Resource Site (insert local and regional newborn screening website information)

Name	
URL	
Comments	

APPENDIX: Resources with Full URL Addresses

Additional Information:

Gene Reviews

<http://www.ncbi.nlm.nih.gov/bookshelf/br.fcgi?book=gene&part=pkupku>

Genetics Home Reference

PKU

<http://ghr.nlm.nih.gov/condition=phenylketonuria>

Tetrahydrobiopterin Deficiency

<http://ghr.nlm.nih.gov/condition=tetrahydrobiopterindeficiency>

Referral (local, state, regional and national):

Testing

http://www.ncbi.nlm.nih.gov/sites/GeneTests/lab/clinical_disease_id/2273?db=genetests&country=United%20States

Clinical Services

<http://www.ncbi.nlm.nih.gov/sites/GeneTests/clinic?db=GeneTests>

Find Genetic Services

<http://www.acmg.net/GIS/Disclaimer.aspx>

Disclaimer: This guideline is designed primarily as an educational resource for clinicians to help them provide quality medical care. It should not be considered inclusive of all proper procedures and tests or exclusive of other procedures and tests that are reasonably directed to obtaining the same results. Adherence to this guideline does not necessarily ensure a successful medical outcome. In determining the propriety of any specific procedure or test, the clinician should apply his or her own professional judgment to the specific clinical circumstances presented by the individual patient or specimen. Clinicians are encouraged to document the reasons for the use of a particular procedure or test, whether or not it is in conformance with this guideline. Clinicians also are advised to take notice of the date this guideline was adopted, and to consider other medical and scientific information that become available after that date.

© American College of Medical Genetics and Genomics, 2012 (Funded in part through MCHB/HRSA/HHS grant #U22MC03957)